Vectors: lines and planes

AS Core Pure Mathematics | Edexcel 8FM0 | 12 questions, 79 marks

Further Maths topic practice

Scan for the solutions

Name: _____ **Date:** _____ **Score:** _____ **/ 79**

1 The points A and B have coordinates (2, −1, 3) and (5, 1, −1) respectively. The line l passes through A and B.

a. Find a vector equation for l.

b. Find a cartesian equation for l.

c. Show that the point C(11, 5, −9) lies on l.

(5 marks)

Answer _____

2 The line l_1 has equation $\mathbf{r} = (\mathbf{i} + 2\mathbf{k}) + \lambda(2\mathbf{i} - \mathbf{j} + 2\mathbf{k})$.

The line l_2 has cartesian equation $\frac{x-3}{4} = \frac{y+1}{-4} = \frac{z}{7}$

Find the acute angle between l_1 and l_2, giving your answer in degrees to one decimal place.

(4 marks)

Answer _____

Free for classroom and home use: print, photocopy and share. Not for resale or re-hosting.

3 The plane Π passes through the point $A(3, -2, 1)$ and is perpendicular to the vector $2\mathbf{i} - \mathbf{j} + 4\mathbf{k}$.

a. Find a vector equation of Π in the form $\mathbf{r} \cdot \mathbf{n} = p$.

b. Write down a cartesian equation of Π.

c. Determine whether the point $B(5, 1, 2)$ lies on Π.

(4 marks)

Answer

4 The lines l_1 and l_2 have equations

l_1: $\mathbf{r} = (2\mathbf{i} + \mathbf{k}) + \lambda(\mathbf{i} + 2\mathbf{j} - \mathbf{k})$

l_2: $\mathbf{r} = (3\mathbf{j} + 4\mathbf{k}) + \mu(\mathbf{i} + \mathbf{j} + 2\mathbf{k})$

a. Show that l_1 and l_2 are skew.

b. The line l_3 has direction $a\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$, where a is a constant. Given that l_3 is perpendicular to l_1, find the value of a.

(6 marks)

Answer

Free for classroom and home use: print, photocopy and share. Not for resale or re-hosting.

5 The lines l_1 and l_2 have equations

l_1: $\mathbf{r} = (\mathbf{i} + 2\mathbf{j} - \mathbf{k}) + \lambda(\mathbf{i} - \mathbf{j} + 2\mathbf{k})$

l_2: $\mathbf{r} = (5\mathbf{i} + \mathbf{j} + 2\mathbf{k}) + \mu(2\mathbf{i} + \mathbf{j} - \mathbf{k})$

a. Show that l_1 and l_2 intersect, and find the coordinates of their point of intersection.

b. Find the acute angle between l_1 and l_2, giving your answer in degrees to one decimal place.

(7 marks)

Answer _____

6 The points $A(1, 0, 2)$, $B(3, 1, 0)$ and $C(0, 2, 1)$ lie in the plane Π.

a. Show that the vector $3\mathbf{i} + 4\mathbf{j} + 5\mathbf{k}$ is perpendicular to both $\overrightarrow{AB}$ and $\overrightarrow{AC}$.

b. Find a cartesian equation of Π.

c. Find the exact perpendicular distance from the origin to Π, giving your answer as a simplified surd.

(7 marks)

Answer _____

Free for classroom and home use: print, photocopy and share. Not for resale or re-hosting.

7 The plane Π has equation $x - 2y + 2z = 9$

The line l has equation $\mathbf{r} = (\mathbf{i} + 3\mathbf{j} - 2\mathbf{k}) + \lambda(2\mathbf{i} + \mathbf{j} + 3\mathbf{k})$.

a. Find the coordinates of the point where l meets Π.

b. Find the acute angle between l and Π, giving your answer in degrees to one decimal place.

(7 marks)

Answer

8 The plane Π_1 has vector equation

$\mathbf{r} = (2\mathbf{i} + \mathbf{j}) + s(\mathbf{i} + 3\mathbf{k}) + t(2\mathbf{j} - \mathbf{k})$

where s and t are scalar parameters.

a. Show that the vector $6\mathbf{i} - \mathbf{j} - 2\mathbf{k}$ is a normal to Π_1.

b. Hence find a cartesian equation of Π_1.

The plane Π_2 has equation $x + y + z = 5$

c. Find the acute angle between Π_1 and Π_2, giving your answer in degrees to one decimal place.

(7 marks)

Answer

Free for classroom and home use: print, photocopy and share. Not for resale or re-hosting.

9 The plane Π has equation $2x + y - 2z = 4$ and the point P has coordinates $(6, 4, -3)$.

a. Find the perpendicular distance from P to Π.

b. Find the coordinates of the foot of the perpendicular from P to Π.

c. The point Q is the reflection of P in Π. Find the coordinates of Q.

(7 marks)

Answer

10 The line l has equation $\mathbf{r} = (\mathbf{i} - \mathbf{j} + 2\mathbf{k}) + \lambda(2\mathbf{i} + \mathbf{j} - 2\mathbf{k})$ and the point A has coordinates $(6, 4, 5)$.

The point F lies on l and AF is perpendicular to l.

a. Find the coordinates of F.

b. Hence find the exact shortest distance from A to l.

The point B on l has parameter $\lambda = 4$

c. Find the exact area of triangle ABF.

(8 marks)

Answer

Free for classroom and home use: print, photocopy and share. Not for resale or re-hosting.

11 A drone is launched from a point A on level ground. Relative to a fixed origin, A has position vector $\mathbf{i} + 2\mathbf{j}$, where $\mathbf{i}$ and $\mathbf{j}$ are horizontal unit vectors due east and due north and $\mathbf{k}$ is a unit vector vertically upwards. Distances are in metres.

The drone flies in a straight line in the direction $2\mathbf{i} + \mathbf{j} + 2\mathbf{k}$ until it reaches the sloping roof of a barn. The roof is modelled as part of the plane with equation $x + 2y + 2z = 29$

a. Find the position vector of the point where the drone reaches the roof.

b. Find the distance the drone travels.

c. Find the acute angle between the path of the drone and the roof, in degrees to one decimal place.

d. Find the shortest distance from A to the plane of the roof.

e. State one limitation of the model.

(8 marks)

Answer

Free for classroom and home use: print, photocopy and share. Not for resale or re-hosting.

12 The line l_1 has equation $\mathbf{r} = (\mathbf{i} + 2\mathbf{j}) + \lambda(\mathbf{i} - \mathbf{j} + 2\mathbf{k})$ and the point P has coordinates (2, 4, 3).

The plane Π contains l_1 and the point P.

a. Show that $7\mathbf{i} + \mathbf{j} - 3\mathbf{k}$ is a normal to Π.

b. Find a cartesian equation of Π.

The line l_2 has equation $\mathbf{r} = (\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}) + \mu(\mathbf{i} + \mathbf{j} + \mathbf{k})$.

c. Find the coordinates of the point where l_2 meets Π.

d. Find the acute angle between l_2 and Π, giving your answer in degrees to one decimal place.

(9 marks)

Answer

End of questions | Total 79 marks

Free for classroom and home use: print, photocopy and share. Not for resale or re-hosting.