Second order non-homogeneous differential equations

A-level Core Pure Mathematics | Edexcel 9FM0 | 12 questions, 95 marks

Further Maths topic practice

Scan for the solutions

Name: _____ **Date:** _____ **Score:** _____ **/ 95**

1 Find the general solution of the differential equation

$$\frac{d^2y}{dx^2} - 3\frac{dy}{dx} + 2y = 4x$$

(4 marks)

Answer _____

2 Find the general solution of the differential equation

$$\frac{d^2y}{dx^2} + 4y = 15 \sin x$$

(4 marks)

Answer _____

3 Find the general solution of the differential equation

$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} - 3y = 10e^{2x}$$

(5 marks)

Answer _____

Free for classroom and home use: print, photocopy and share. Not for resale or re-hosting.

4 The differential equation

$$\frac{d^2y}{dx^2} - \frac{dy}{dx} - 6y = 10e^{3x}$$

is to be solved.

a. Find the complementary function and explain why a particular integral of the form $y = \lambda e^{3x}$ cannot be used.

b. Using a particular integral of the form $y = \lambda x e^{3x}$, find the general solution of the differential equation.

(6 marks)

Answer

5 The differential equation

$$\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 4y = 8x^2$$

is to be solved.

a. Find the general solution of the differential equation.

b. Find the particular solution for which $y = 1$ and $\frac{dy}{dx} = 2$ when $x = 0$

(8 marks)

Answer

Free for classroom and home use: print, photocopy and share. Not for resale or re-hosting.

6 The differential equation

$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + 5y = 10 \sin x$$

is to be solved.

a. Find a particular integral of the differential equation.

b. Find the particular solution for which $y = 0$ and $\frac{dy}{dx} = 0$ when $x = 0$

c. Describe the behaviour of y for large values of x, stating the amplitude of the oscillation as an exact value.

(9 marks)

Answer

Free for classroom and home use: print, photocopy and share. Not for resale or re-hosting.

7 The differential equation

$$\frac{d^2y}{dx^2} + 16y = 24 \cos 4x$$

is to be solved.

a. Show that a particular integral of the differential equation is $y = 3x \sin 4x$

b. Hence find the general solution of the differential equation.

c. Find the particular solution for which $y = 2$ and $\frac{dy}{dx} = 8$ when $x = 0$

d. Find the exact value of y when $x = \frac{\pi}{8}$ for this particular solution.

(9 marks)

Answer

Free for classroom and home use: print, photocopy and share. Not for resale or re-hosting.

8 The differential equation

$$\frac{d^2y}{dx^2} - 6\frac{dy}{dx} + 9y = 6e^{3x}$$

is to be solved.

a. Explain why a particular integral of the form $\lambda x^2 e^{3x}$ is needed, and find the value of λ.

b. Find the particular solution for which $y = 1$ and $\frac{dy}{dx} = 5$ when $x = 0$

c. Show that $y > 0$ for all values of x.

(8 marks)

Answer

9 The differential equation

$$\frac{d^2y}{dx^2} + \frac{dy}{dx} - 6y = 13 - 6x + 12e^{-x}$$

is to be solved.

a. Find the general solution of the differential equation.

A particular solution has $y = 0$ when $x = 0$, and for this solution $y - x$ remains finite as $x \to \infty$

b. Find this particular solution.

c. Find the value of $\frac{dy}{dx}$ when $x = 0$ for this particular solution.

(9 marks)

Answer

10 The temperature, θ °C, inside an industrial oven t minutes after it is switched on is modelled by the differential equation

$$\frac{d^2\theta}{dt^2} + 2\frac{d\theta}{dt} + 10\theta = 1800$$

When the oven is switched on, its temperature is 20 °C and is not changing.

a. Find the general solution of the differential equation.

b. Show that the particular solution is

$\theta = 180 - e^{-t}(160 \cos 3t + \frac{160}{3} \sin 3t)$

c. Show that $\frac{d\theta}{dt} = \frac{1600}{3} e^{-t} \sin 3t$

d. Hence find, according to the model, the maximum temperature in the oven, giving your answer to the nearest degree.

e. State the temperature that the oven settles at in the long term, according to the model.

(11 marks)

Answer _____

Free for classroom and home use: print, photocopy and share. Not for resale or re-hosting.

11 A sledge is pulled from rest across horizontal snow. Its displacement, x metres, from its starting point t seconds after it starts to move is modelled by the differential equation

$$\frac{d^2x}{dt^2} + 4\frac{dx}{dt} = 8$$

a. Find the complementary function and explain why a particular integral of the form $x = k$, where k is a constant, will not work.

b. Using a particular integral of the form $x = \lambda t$, find the particular solution for the motion of the sledge.

c. Find, according to the model, the distance travelled by the sledge in the first 3 seconds.

d. Find, according to the model, the speed of the sledge after 0.5 seconds, and state the speed that the sledge approaches.

(10 marks)

Answer

Free for classroom and home use: print, photocopy and share. Not for resale or re-hosting.

12 A particle P moves along the x-axis. At time t seconds, $t \geq 0$, P is x metres from the origin O, where

$$\frac{d^2x}{dt^2} + 3\frac{dx}{dt} + 2x = 2t + 7$$

a. Find the general solution of the differential equation.

b. Find the particular solution for which $x = 5$ and $\frac{dx}{dt} = -4$ when $t = 0$

c. Show that the particle is momentarily at rest when $e^{-t} = \frac{\sqrt{17} - 1}{8}$

d. Find, according to the model, the minimum distance between P and O, giving your answer to 3 significant figures, and justify that it is a minimum.

e. Describe the motion of P for large values of t.

(12 marks)

Answer

End of questions | Total 95 marks

Free for classroom and home use: print, photocopy and share. Not for resale or re-hosting.