



- 1 In triangle ABC, $AB = 3$ cm, $AC = 5$ cm and $BC = \sqrt{19}$ cm.
Find the size of angle BAC.




Diagram NOT accurately drawn

(4 marks)

Angle BAC is opposite BC, so use the cosine rule for the angle

$$\cos A = \frac{AB^2 + AC^2 - BC^2}{2 \times AB \times AC}$$

$$\cos A = \frac{9 + 25 - 19}{2 \times 3 \times 5} = \frac{15}{30} = \frac{1}{2}$$

$$\cos A = \frac{1}{2}, \text{ so angle BAC} = 60^\circ$$

$$\boxed{60^\circ}$$

- 2 Show that $\sqrt{12} + \sqrt{27}$ can be written as $5\sqrt{3}$.

(2 marks)

$$\sqrt{12} = \sqrt{4 \times 3} = 2\sqrt{3}$$

$$\sqrt{27} = \sqrt{9 \times 3} = 3\sqrt{3}$$

$$2\sqrt{3} + 3\sqrt{3} = 5\sqrt{3}$$

$$\boxed{5\sqrt{3}}$$

- 3 ABCD is a parallelogram. AB = 12 cm, AD = 7.5 cm and angle DAB = 65° .
Calculate the area of the parallelogram. Give your answer correct to 3 significant figures.

(2 marks)

Parallelogram = 2 identical triangles

$$\text{Area} = 2 \times \frac{1}{2} \times 12 \times 7.5 \times \sin 65^\circ$$

$$\text{Area} = 12 \times 7.5 \times \sin 65^\circ$$

$$\text{Area} = 81.568...$$

$$\boxed{81.6 \text{ cm}^2}$$

- 4 A and B are two events.
 $P(A) = 0.5$, $P(B) = 0.4$ and $P(A \cup B) = 0.7$

a. Work out $P(A \cap B)$.

b. Work out $P(B | A)$.

(4 marks)

a)

$$0.7 = 0.5 + 0.4 - P(A \cap B)$$

$$P(A \cap B) = 0.9 - 0.7$$

$$P(A \cap B) = 0.2$$

b)

$$P(B | A) = \frac{P(A \cap B)}{P(A)}$$

$$P(B | A) = \frac{0.2}{0.5}$$

$$\boxed{\text{a) } 0.2 \text{ b) } 0.4}$$

5

A, B and C are points on a circle with centre O.

B lies on the major arc AC.

Angle AOC = 128° .

Work out the size of angle ABC. Give a reason for your answer.




Diagram NOT accurately drawn

(3 marks)

The angle at the centre of a circle is twice the angle at the circumference when both stand on the same arc

$$\text{Angle } ABC = 128^\circ \div 2 = 64^\circ$$

64

6

The line $y = ax + 5$ is perpendicular to the line $by = 3x - 2$.

Find b in terms of a .

(3 marks)

$$m_1 = a$$

$$y = \frac{3}{b}x - \frac{2}{b}$$

$$m_2 = \frac{3}{b}$$

Perpendicular: product of gradients = -1

$$a \times \frac{3}{b} = -1$$

$$\begin{array}{ccc} \text{xb} & 3a & = -b \quad \text{xb} \\ & \parallel & \\ & 3a & = -b \end{array}$$

$$b = -3a$$

7

Work out the value of $(\frac{36}{25})^{\frac{1}{2}}$

(2 marks)

A power of $\frac{1}{2}$ means the square root of the numerator and of the denominator

$$\sqrt{36} = 6 \text{ and } \sqrt{25} = 5, \text{ so the answer is } \frac{6}{5}$$

$$\boxed{\frac{6}{5}}$$

8

Write $x^2 + 8x + 3$ in the form $(x + a)^2 + b$, where a and b are integers.

(2 marks)

Expanding $(x + a)^2$ gives $x^2 + 2ax + a^2$. To match $8x$, we need $2a = 8$, so $a = 4$

Halve the coefficient of x

$$(x + 4)^2 = x^2 + 8x + 16$$

Adjust the constant

$$x^2 + 8x + 3 = (x + 4)^2 - 16 + 3 = (x + 4)^2 - 13$$

Check by expanding

$$x^2 + 8x + 16 - 13 = x^2 + 8x + 3$$

$$\boxed{(x + 4)^2 - 13}$$

9

Two mathematically similar candles, C and D, are made from the same wax.

C has surface area 36 cm^2 and D has surface area 81 cm^2 .The volume of C is 24 cm^3 .

Work out the volume of D.

(3 marks)

Area ratio C

$$0 = 36 : 81 = 4 : 9$$

$$\text{Length ratio} = \sqrt{4} : \sqrt{9} = 2 : 3$$

$$\text{Volume ratio} = 2^3 : 3^3 = 8 : 27$$

$$\text{Volume of D} = 24 \times 27 \div 8 = 81 \text{ cm}^3$$

$$\boxed{81 \text{ cm}^3}$$

10

The frequency densities for the times, t minutes, that 80 customers waited in a post office queue are shown in the table.

Time (t minutes)	Frequency density
$0 < t \leq 10$	2
$10 < t \leq 30$	1.5
$30 < t \leq 40$	3

Mia says, "The $0 < t \leq 10$ class must have the most customers because its bar is tall." Is Mia correct? You must show your working.

(2 marks)

Frequency = frequency density $\times$ class width

$$0-10: 2 \times 10 = 20$$

$$10-30: 1.5 \times 20 = 30$$

$$30-40: 3 \times 10 = 30$$

No - 0-10 has the fewest (20)

11

Factorise $2x^2 + 7x + 3$

(2 marks)

Look for two numbers that multiply to $2 \times 3 = 6$ and add to 7: they are 6 and 1

Split the middle term

$$2x^2 + 6x + x + 3 = 2x(x + 3) + 1(x + 3)$$

Both groups now contain the whole factor $(x + 3)$. Taking out that shared factor leaves $2x + 1$, giving the two brackets below

Factorise: $(2x + 1)(x + 3)$

Check by expanding

$$2x^2 + 6x + x + 3 = 2x^2 + 7x + 3, \text{ correct}$$

$(2x + 1)(x + 3)$

- 12 In triangle ABC, angle BAC = 52° , angle ACB = 63° and AC = 11 cm.
Calculate the length of BC. Give your answer correct to 3 significant figures.




Diagram NOT accurately drawn

(3 marks)

$$\text{Angle } ABC = 180^\circ - 52^\circ - 63^\circ = 65^\circ$$

BC is opposite angle A and AC is opposite angle B

Sine rule

$$\frac{BC}{\sin A} = \frac{AC}{\sin B}$$

$$BC = 11 \times \sin 52^\circ \div \sin 65^\circ = 11 \times 0.78801... \div 0.90630..$$

$$BC = 9.564... = 9.56 \text{ cm (3 s.f.)}$$

$$\boxed{9.56 \text{ cm}}$$

- 13 p is inversely proportional to q .

When $q = 5$, $p = 12$.

- Find a formula for p in terms of q .
- Find the value of p when $q = 8$.

(3 marks)

Inversely proportional means that the product pq is constant

$$pq = k. \text{ Dividing by } q \text{ gives } p = k/q$$

Substitute

$$12 = \frac{k}{5}, \text{ so } k = 60 \text{ and } p = \frac{60}{q}$$

$$\text{When } q = 8, p = 60 \div 8 = 7.5$$

- 14 A park keeper wants to estimate the number of fish in a lake.
She catches 30 fish, tags them and puts them back in the lake.
The next day she catches 40 fish. 8 of these fish are tagged.
Work out an estimate for the number of fish in the lake.

(2 marks)

$$\text{Proportion of tagged fish in the second sample} = \frac{8}{40} = \frac{1}{5}$$

Assume the same proportion of the whole population is tagged, so 30 fish is $\frac{1}{5}$ of the population

$$\text{Estimate} = 30 \times 40 \div 8 = 150$$

150

- 15 $f(x) = 3x - 4$ $g(x) = x^2$

a. Find $f(5)$

b. Find $gf(2)$

(3 marks)

$gf(x)$ means $g(f(x))$: apply f first, then put its output into g . The function nearest the input is done first; gf does not mean multiply g by f

$$\text{a) } f(5) = 3(5) - 4 = 11$$

$$\text{b) } f(2) = 3(2) - 4 = 2, \text{ then } g(2) = 2^2 = 4$$

- 16 Here are four equations.

$$\text{A: } y = 3 \times 1.2^x \quad \text{B: } y = 5 \times 0.8^x \quad \text{C: } y = 0.5 \times 4^x \quad \text{D: } y = 2 \times \left(\frac{1}{3}\right)^x$$

Which of the graphs show exponential decay (the curve goes down as x increases)? Give a reason.

(2 marks)

$$y = a \times b^x$$

$b > 1$ means growth

$0 < b < 1$ means decay

A: $b = 1.2$ growth

B: $b = 0.8$ decay

C: $b = 4$ growth

D: $b = \frac{1}{3}$ decay

B and D

- 17 A histogram shows the heights of 71 trees. The table gives the frequency density for each class.

Height (h m)	Frequency density
$0 < h \leq 10$	1.2
$10 < h \leq 20$	2.5
$20 < h \leq 40$	1.5
$40 < h \leq 50$	0.4

Work out the frequency for each class and check that the total is 71.

(2 marks)

$$\text{frequency} = f.d \times \text{width}$$

$$0-10: 1.2 \times 10 = 12$$

$$10-20: 2.5 \times 10 = 25$$

$$20-40: 1.5 \times 20 = 30$$

$$40-50: 0.4 \times 10 = 4$$

$$12 + 25 + 30 + 4 = 71$$

$$12, 25, 30, 4$$

- 18 Here are the first five terms of a quadratic sequence.

2 7 14 23 34

Find an expression, in terms of n , for the n th term of this sequence.

(2 marks)

For a quadratic sequence $an^2 + bn + c$, the second difference is $2a$. The n^2 terms 1, 4, 9, 16, ... have first differences 3, 5, 7, ... and second difference 2; multiplying by a makes it $2a$. The linear part $bn + c$ adds no second difference

First differences

5, 7, 9, 11. Second difference: 2, so the n^2 coefficient is 2 divided by 2 = 1

Subtract n^2 (1, 4, 9, 16, 25) from the sequence: 1, 3, 5, 7, 9, which has n th term $2n - 1$

So the n th term is $n^2 + 2n - 1$

Check

$n = 1$ gives $1 + 2 - 1 = 2$; $n = 2$ gives $4 + 4 - 1 = 7$; $n = 3$ gives $9 + 6 - 1 = 14$; $n = 4$ gives $16 + 8 - 1 = 23$; $n = 5$ gives $25 + 10 - 1 = 34$

$$n^2 + 2n - 1$$

- 19 Prove algebraically that the sum of an odd number and an even number is always odd.

(2 marks)

Even number: $2n$

Odd number: $2m + 1$ (n, m integers)

$$\begin{aligned} 2n + (2m + 1) &= 2n + 2m + 1 \\ &= 2(n + m) + 1 \end{aligned}$$

$2(n + m)$ is a multiple of 2, so it is even; one more makes it odd

Sum is always odd

20

Simplify fully $\frac{2x^2 + 6x}{4x}$

(2 marks)

Factorise the top

$$\begin{aligned} 2x^2 + 6x &= 2x(x + 3) \\ &= \frac{2x(x + 3)}{4x} \end{aligned}$$

Cancel the common factor $2x$

$$\boxed{\frac{x + 3}{2}}$$