



- 1 Solve the simultaneous equations

$$x^2 + y^2 = 20$$

$$y = 2x$$

(4 marks)

Substitute $y = 2x$ into the circle equation: $x^2 + 4x^2 = 20$

So $5x^2 = 20$, giving $x^2 = 4$

$$x = 2 \text{ or } x = -2$$

When $x = 2$, $y = 4$. When $x = -2$, $y = -4$

Check

$$4 + 16 = 20 \text{ for both pairs}$$

$$x = 2, y = 4 \text{ and } x = -2, y = -4$$

2

OAB is a triangle. $OA = \mathbf{a}$ and $OB = \mathbf{b}$.

M is the midpoint of OA and N is the midpoint of OB.

Show that MN is parallel to AB.




Diagram NOT accurately drawn

(3 marks)

$$OM = \frac{1}{2}\mathbf{a} \text{ and } ON = \frac{1}{2}\mathbf{b}$$

$$MN = ON - OM = \frac{1}{2}\mathbf{b} - \frac{1}{2}\mathbf{a} = \frac{1}{2}(\mathbf{b} - \mathbf{a})$$

$$AB = OB - OA = \mathbf{b} - \mathbf{a}$$

$MN = \frac{1}{2}AB$, so MN is a scalar multiple of AB and is therefore parallel to AB

A scalar is an ordinary number multiplying a vector. Multiplying a non-zero vector by $1/2$ halves its length without changing its direction, which establishes parallel lines

3

The curve $\mathbf{C}$ has equation $y = \cos x^\circ$ for $0 \leq x \leq 360$.

Write down the coordinates of the minimum point of $\mathbf{C}$.

(2 marks)

The cosine curve starts at $(0, 1)$, falls to its lowest value of -1 at $x = 180$, and returns to 1 at $x = 360$

So the minimum point is $(180, -1)$

$(180, -1)$

4 OAB is a triangle.

Vector $OA = \mathbf{a}$ and vector $OB = \mathbf{b}$.

M is the midpoint of OA and N is the midpoint of OB.

Prove that MN is parallel to AB.




Diagram NOT accurately drawn

(4 marks)

Vector $OM = \frac{1}{2}\mathbf{a}$ because M is the midpoint of OA

Vector $ON = \frac{1}{2}\mathbf{b}$ because N is the midpoint of OB

Vector $MN = ON - OM = \frac{1}{2}\mathbf{b} - \frac{1}{2}\mathbf{a} = \frac{1}{2}(\mathbf{b} - \mathbf{a})$

Vector $AB = OB - OA = \mathbf{b} - \mathbf{a}$

A scalar is an ordinary number. Multiplying a nonzero vector by a scalar changes its length (and may reverse it), but keeps its direction parallel

So $MN = \frac{1}{2}AB$, which means MN is a scalar multiple of AB

Therefore MN is parallel to AB

5 Expand and simplify $(5 + 2\sqrt{3})(5 - 2\sqrt{3})$

(2 marks)

difference of two squares

$$\begin{aligned}(5 + 2\sqrt{3})(5 - 2\sqrt{3}) &= 5^2 - (2\sqrt{3})^2 \\ &= 25 - 4 \times 3 \\ &= 25 - 12\end{aligned}$$

6

The graph of $y = \sin x$ for $0^\circ \leq x \leq 360^\circ$ has a maximum point at $(90, 1)$.

a. Write down the coordinates of the maximum point of the graph of $y = \sin x - 3$

b. Write down the coordinates of the maximum point of the graph of $y = \sin(x - 30^\circ)$

(2 marks)

a) $y = \sin x - 3$ translates the graph 3 units down, so $(90, 1)$ maps to $(90, -2)$

b) $y = \sin(x - 30)$ translates the graph 30 degrees to the right, so $(90, 1)$ maps to $(120, 1)$

7

Solve $(x - 3)(x + 7) \geq 0$

(2 marks)

The roots of $(x - 3)(x + 7) = 0$ are $x = 3$ and $x = -7$

The parabola is U-shaped, so the product is greater than or equal to zero outside the roots (including the roots themselves because the inequality is not strict)

So $x = 3$

$x = 3$

8

The number of members of a gym is modelled by the rule

$$u_{n+1} = 1.2u_n - 50$$

where u_n is the number of members after n years.

There are 500 members now, so $u_0 = 500$.

Work out the number of members the model predicts after 2 years.

(2 marks)

u_0 is the starting value. The subscript $n + 1$ means the next year: put the current value u_n into the rule to get the next value. Repeat the rule twice for two years

$$u_1 = 1.2 \times 500 - 50 = 600 - 50 = 550$$

$$u_2 = 1.2 \times 550 - 50 = 660 - 50 = 610$$

610

9

A runner's velocity-time graph is a smooth curve. The table gives her velocity v m/s at time t seconds.

t : 0, 10, 20, 30 v : 0, 4.2, 6.6, 7.8

Use the trapezium rule with 3 strips of equal width to estimate the distance the runner travels in the first 30 seconds.

(3 marks)

Distance is the area under the velocity-time graph. Use 3 trapezia of width 10

First strip

$$\frac{1}{2} \times 10 \times (0 + 4.2) = 21$$

Second strip

$$\frac{1}{2} \times 10 \times (4.2 + 6.6) = 54$$

Third strip

$$\frac{1}{2} \times 10 \times (6.6 + 7.8) = 72$$

$$\text{Estimate} = 21 + 54 + 72 = 147 \text{ m}$$

147 m

10

ABCDEFGH is a cuboid with base ABCD and top face EFGH, where E is directly above A, F is directly above B, G is directly above C and H is directly above D.

AB = 3 cm, BC = 4 cm and CG = 12 cm.

Work out the length of AG.




Diagram NOT accurately drawn

(2 marks)

First find the diagonal AC of the base using Pythagoras in triangle ABC: $AC^2 = 3^2 + 4^2 = 9 + 16 = 25$, so $AC = 5$ cm

AG is the space diagonal. Triangle ACG has a right angle at C because CG is vertical

$$AG^2 = AC^2 + CG^2 = 25 + 144 = 169$$

$$AG = \sqrt{169} = 13 \text{ cm}$$

Alternatively, in one step

$$AG = \sqrt{(3^2 + 4^2 + 12^2)} = \sqrt{169} = 13 \text{ cm}$$

13 cm

11

A circle has its centre at the origin O. The diameter of the circle is 16.

Find the equation of the circle.

(2 marks)

$$\text{radius} = 16 \div 2 = 8$$

$$r^2 = 8^2 = 64$$

$$x^2 + y^2 = 64$$

12 $f(x) = x^3 - 4x - 3$

Show that the equation $f(x) = 0$ has a root between $x = 2$ and $x = 3$.

(2 marks)

$$f(2) = 8 - 8 - 3 = -3$$

$$f(3) = 27 - 12 - 3 = 12$$

Sign change and f is continuous

A root lies between $x = 2$ and $x = 3$

13 Write $\frac{x+1}{2} + \frac{2x-1}{5}$ as a single fraction in its simplest form.

(2 marks)

$$\begin{aligned} \frac{x+1}{2} + \frac{2x-1}{5} &= \frac{5(x+1)}{10} + \frac{2(2x-1)}{10} \\ &= \frac{5x+5+4x-2}{10} \end{aligned}$$

$$\frac{9x+3}{10}$$

14 The temperature, T °C, of a cup of tea is plotted against time, t minutes, and the points form a smooth curve going down.

At $t = 10$ a tangent is drawn to the curve. The tangent passes through the points (2, 80) and (14, 56).

- Work out the gradient of the tangent.
- What does this gradient tell you about the tea at $t = 10$?

(3 marks)

a)

$$\begin{aligned} \text{gradient} &= \{56 - 80\} / \{14 - 2\} \\ &= \{-24\} / \{12\} \\ &= -2 \end{aligned}$$

b)

gradient = rate of change of T
negative: temperature is falling

a) -2 b) cooling at 2 °C per minute

15

The curve **C** has equation $y = x^2 - 4x + 7$

The line **L** has equation $y = 2x - 2$

Show that **L** touches **C** at exactly one point, and find the coordinates of this point.

(4 marks)

Set the expressions for y equal

$$x^2 - 4x + 7 = 2x - 2$$

Rearrange

$$x^2 - 6x + 9 = 0$$

Factorise

$$(x - 3)^2 = 0$$

Both factors in $(x - 3)(x - 3)$ are zero at the same value, $x = 3$; this is called a repeated root. Also, curve minus line $= (x - 3)^2$ is never negative, so the curve stays on one side of the line and touches it at this one point


When $x = 3$, $y = 2(3) - 2 = 4$, so the point is $(3, 4)$

Check on the curve

$$9 - 12 + 7 = 4$$

16 Sketch the graph of $y = \tan x^\circ$ for $0 \leq x \leq 360$.

Show clearly where the graph crosses the x -axis and mark the values of x where the graph has asymptotes.



(2 marks)

The tangent graph crosses the x -axis at $x = 0, 180$ and 360

It has vertical asymptotes at $x = 90$ and $x = 270$

Each branch increases from bottom left to top right, repeating every 180

A vertical asymptote is a line that the graph approaches while its values grow without bound. Here $\tan x = \sin x \div \cos x$ is undefined at 90° and 270° because $\cos x = 0$. Do not join the branches across those lines

17 The graph of $y = f(x)$ has a turning point at $(3, -5)$.

a. Write down the coordinates of the turning point of the graph of $y = f(x) + 4$

b. Write down the coordinates of the turning point of the graph of $y = f(x - 2)$

(2 marks)

a) $y = f(x) + 4$ translates the graph 4 units up, so $(3, -5)$ maps to $(3, -5 + 4) = (3, -1)$

b) $y = f(x - 2)$ translates the graph 2 units to the right, so $(3, -5)$ maps to $(3 + 2, -5) = (5, -5)$

For part (b), the original turning point occurs when the input of f is 3. In $f(x - 2)$, solve $x - 2 = 3$, giving $x = 5$. This explains why subtracting inside the function moves the point to the right

18

The circle C has equation $x^2 + y^2 = 61$

a. Show that the point A (5, -6) lies on C.

b. Does the point B (4, 7) lie on C? You must justify your answer.

(3 marks)

a) Substitute A into the left-hand side

$$5^2 + (-6)^2 = 25 + 36 = 61. \text{ This equals 61, so A lies on C}$$

b) Substitute B

$$4^2 + 7^2 = 16 + 49 = 65. \text{ Since 65 is not equal to 61, B does not lie on C. In fact } 65 > 61, \text{ so B is outside the circle}$$

19

Erin is asked to rationalise the denominator of $\frac{8}{\sqrt{6}}$

Here is her working.

$$\frac{8}{\sqrt{6}} = \frac{8 \times \sqrt{6}}{\sqrt{6} \times \sqrt{6}} = \frac{8\sqrt{6}}{36}$$

Erin has made one mistake.

Explain the mistake Erin has made.

(2 marks)

Erin has squared 6 instead of squaring $\sqrt{6}$

$$\sqrt{6} \times \sqrt{6} = 6, \text{ not } 36$$

The correct answer is $\frac{8\sqrt{6}}{6}$, which simplifies to $\frac{4\sqrt{6}}{3}$

20

Rationalise the denominator of $\frac{12}{\sqrt{3}}$

Give your answer in its simplest form.

(2 marks)

Rationalise means remove the square root from the denominator. Multiplying top and bottom by $\sqrt{3}$ multiplies the fraction by $\sqrt{3}/\sqrt{3} = 1$, so it keeps the same value; $\sqrt{3} \times \sqrt{3} = 3$

Multiply the numerator and denominator by $\sqrt{3}$

$$\frac{12}{\sqrt{3}} = \frac{12\sqrt{3}}{3}$$

Divide top and bottom by 3

$$\frac{12\sqrt{3}}{3} = 4\sqrt{3}$$

$$\boxed{4\sqrt{3}}$$